THE LANCET Child & Adolescent Health

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Kikkenborg Berg S, Nielsen SD, Nygaard U, et al. Long COVID symptoms in SARS-CoV-2-positive adolescents and matched controls (LongCOVIDKidsDK): a national, cross-sectional study. *Lancet Child Adolesc Health* 2022; published online Feb 7. https://doi.org/10.1016/S2352-4642(22)00004-9.

Contents supplementary

Investigators	2
Table S1- Duration of all symptoms in cases and controls.	3
Table S2 - Psychological and social symptoms in cases and controls	6
Table S3 - Sick days during the past 12 months	7
Table S4 - Questionnaire LongCOVIDKidsDK	8
Table S5 - Covid-19 Denmark: School/national lockdown and reopening, and changes in testing	
policies	10

Investigators

Selina Kikkenborg Berg, RN, MScN, Ph.D. a,b, Susanne Dam Nielsen MD, PhDb,d, Ulrikka Nygaard MD, PhDb,c,

Henning Bundgaard, MD, PhDa, Pernille Palm, RN, MScN, PhDa, Camilla Rotvig, RN, MPHa, Anne Vinggaard

Christensen, MSPH, PhDa

^aDepartment of Cardiology, Rigshospitalet, Copenhagen University Hospital, Inge Lehmanns Vej 7, 2100 Copenhagen,

Denmark

^bFaculty of Health and Medical Sciences, University of Copenhagen, Blegdamsvej 3B, 2200 Copenhagen N, Denmark

^cDepartment of Paediatrics and Adolescents medicine, Rigshospitalet, Copenhagen University Hospital, Blegdamsvej 9,

2100 Copenhagen, Denmark

^dDepartment of Infectious Disease, Rigshospitalet, Copenhagen University Hospital, Blegdamsvej 9, 2100 Copenhagen,

Denmark

Corresponding author:

Professor Selina Kikkenborg Berg

Rigshospitalet

The Heart Centre, Unit 2151

Blegdamsvej 9

2100 Copenhagen

Denmark

Phone: +45 35459526

E-mail: selina@rh.dk

2

Table S1- Duration of all symptoms in cases and controls.

	At least 2 months	At least 3 months	At least 6 months	At least 9 months	At least 12 months
	Cases/controls	Cases/controls	Cases/controls	Cases/controls	Cases/controls
Total n	5106/21640	5106/21640	4250/16257	1085/10852	242/5261
Stomach aches, n (%)					
Almost never and Sometimes	240 (4·7)/ 822 (3·8)	212 (4·2)/ 706 (3·3)	135 (3·2)/ 414 (2·5)	28 (2·6)/ 227 (2·1)	7 (2·9)/ 115 (2·2)
Often and Almost always	119 (2·3)/ 626 (2·9)*a	113 (2·2)/ 576 (2·7)	74 (1·7)/ 367 (2·3)*	9 (0·8)/ 196 (1·8)	0 (0·0)/ 87 (1·7)
Chest pain, n (%)	020 (2))	070(27)	201 (20)	170 (1 0)	0, (1, 1)
Almost never and Sometimes	382 (7·5)/ 1083 (5·0)	333 (6·5)/ 938 (4·3)	231 (5·4)/ 583 (3·6)	51 (4·7)/ 324 (3·0)	7 (2·9)/ 153 (2·9)
Often and Almost always	125 (2·5)/ 384 (1·8)*	115 (2·3)/ 335 (1·5)	77 (1·8)/ 216 (1·3)*	13 (1·2)/ 116 (1·1)	1-4/ 48 (0·9)
Headache, n (%)	304 (1 0)	333 (1 3)	210 (1 3)	110 (1 1)	40 (0 2)
Almost never and Sometimes	728 (14·3)/ 1835 (8·5)	648 (127)/ 1589 (7·3)	454 (10·7)/ 1029 (6·3)	96 (8·8)/ 588 (5·4)	17 (7·0)/ 292 (5·6)
Often and Almost always	610 (12·0)/ 2166 (10·0)***	567 (11·1)/ 1929 (8·9)	1029 (0·3) 419 (9·9)/ 1261 (7·8)***	87 (8·0)/ 694 (6·4)	15 (6·2)/ 309 (5·9)
Fatigue, n (%)	2100 (10-0)	1929 (8.9)	1201 (7.8)	094 (0.4)	309 (3.9)
Almost never and Sometimes	781 (15·3)/ 2381 (11·0)	692 (13·6)/ 1904 (8·8)	465 (10·9)/ 1081 (6·6)	90 (8·3)/ 564 (5·2)	11 (4·5)/ 234 (4·4)
Often and Almost always	1057 (20·7)/ 4249 (19·6)	979 (19·2)/ 3810 (17·6)	694 (16·3)/ 2395 (14·7)*	144 (13·3)/ 1296 (11·9)	19 (7·9)/ 542 (10·3)
Pain in muscles/joints, n (%)	1219 (19 0)	3010 (17 0)	2333 (117)	1250 (11 5)	3 12 (10 3)
Almost never and Sometimes	351 (6·9)/ 980 (4·5)	293 (5·7)/ 816 (3·8)	184 (4·3)/ 461 (2·8)	41 (3·8)/ 257 (2·4)	7 (2·9)/ 116 (2·2)
Often and Almost always	163 (3·2)/ 633 (2·9)	157 (3·1)/ 587 (2·7)	110 (2·6)/ 388 (2·4)	23 (2·1)/ 23 (2·0)	5 (2·1)/ 100 (1·9)
Sore throat, n (%)	033 (2.9)	387 (2.7)	388 (2.4)	219 (2.0)	100 (1.3)
Almost never and Sometimes	222 (4·3)/	192 (3.8)/	131 (3·1)/	29 (2.7)/	5 (2·1)/ 58 (1·1)
Often and Almost always	509 (2·4) 70 (1·4)/	404 (1·9) 62 (1·2)/	225 (1·4) 39 (0·9)/	128 (1·2) 10 (0·9)/	1-4/
7.	185 (0.9)**	151 (0.7)	80 (0.5)*	41 (0.4)	13 (0.2)
Dizziness , n (%) Almost never and Sometimes	506 (9.9)/	454 (8.9)/	305 (7.2)/	62 (5.7)/	14 (5.8)/
Often and Almost always	1334 (6·2) 220 (4·3)/	1116 (5·2) 194 (3·8)/	647 (4·0) 130 (3·1)/	335 (3·1) 27 (2·5)/	145 (2·8) 1-4/
•	696 (3.2)**	619 (2.9)	386 (2·4)*	200 (1.8)	79 (1.5)
Rashes, n (%)					
Almost never and Sometimes	163 (3·2)/ 645 (3·0)	140 (2·7)/ 502 (2·3)	94 (2·2)/ 258 (1·6)	22 (2·0)/ 146 (1·3)	3 (1·2)/ 67 (1·3)
Often and Almost always	134 (2·6)/ 771 (3·6)**	129 (2·5)/ 716 (3·3)	93 (2·2)/ 507 (3·1)*	26 (2·4)/ 288 (2·7)	1-4/ 139 (2·6)
Mood swings, n (%)	//1 (3.0)	/10 (3.3)	307 (3.1)	200 (2.1)	139 (2.0)

Almost never and Sometimes	619 (12·1)/	563 (11.0)	401 (9.4)/	81 (7.5)/	18 (7.4)/
Annost never and Sometimes	2359 (10.9)	/2034 (9.4)	1206 (7.4)	657 (6.1)	306 (5.8)
Often and Almost always	554 (10.9)/	527 (10·3)/	390 (9·2)/	78 (7.2)/	15 (6·2)/
Often and Annost arways	2756 (12.7)**	2555 (11.8)	1646 (10.1)	918 (8.5)	414 (7.9)
Nausea, n (%)	2/30 (12-7)***	2333 (11-8)	1040 (10-1)	918 (8.3)	414 (7.9)
, (,	244 (6.7)/	200 (5.0)/	100 (4.5)/	25 (2.2)/	9 (2 2)/
Almost never and Sometimes	344 (6.7)/	299 (5.9)/	190 (4.5)/	35 (3·2)/	8 (3·3)/
	1114 (5·1)	893 (4.1)	540 (3·3)	297 (2.7)	139 (2.6)
Often and Almost always	179 (3.5)/	159 (3.1)/	108 (2.5)/	26 (2·4)/	1-4/
	692 (3.2)	625 (2.9)	372 (2·3)	183 (1.7)	1-4
Fever, n (%)					
Almost never and Sometimes	78 (1.5)/	66 (1.3)/	45 (1·1)/	7 (0.6)/	1-4/
	191 (0.9)	162 (0.7)	115 (0.7)	64 (0.6)	36 (0.7)
Often and Almost always	10 (0.2)/	6 (0.1)/	1-4)/	1-4/	0 (0.0)/
	27(0.1)	24 (0·1)	19 (0·1)	8 (0.1)	1-4
Loss of appetite, n (%)					
Almost never and Sometimes	454 (8.9)/	365 (7.1)/	220 (5·2)/	34 (3·1)/	7 (2.9)/
	1443 (6.7)	1128 (5.2)	544 (3.3)	254 (2.3)	115 (2.2)
Often and Almost always	403 (7.9)/	352 (6.9)/	229 (5.4)/	48 (4.4)/	9 (3.7)/
y	1498 (6.9)*	1265 (5.8)	719 (4.4)*	364 (3.4)	153 (2.9)
Trouble breathing, n (%)		, ,			,
Almost never and Sometimes	427 (8.4)/	385 (7.5)/	243 (5.7)/	44 (4·1)/	9 (3.7)/
Timost never and sometimes	796 (3.7)	686 (3.2)	411 (2.5)	228 (2·1)	112 (2.1)
Often and Almost always	273 (5.4)/	254 (5.0)/	177 (4.2)/	39 (3.6)/	5 (2·1)/
Often and Annost arways	444 (2·1)***	416 (1.9)	281 (1.7)***	153 (1.4)	60 (1·1)
Dark circles under the eyes, n (%)	1 444 (2.1)	410 (1-5)	201 (117)	133 (1.4)	00 (1-1)
Almost never and Sometimes	221 (6 2)/	297 (5.8)/	208 (4.9)/	51 (4.7)/	11 (4.5)/
Almost never and Sometimes	321 (6.3)/				11 (4.5)/
00 141 1	1241 (5.7)	1051 (4.9)	590 (3.6)	313 (2.9)	131 (2.5)
Often and Almost always	448 (8.8)/	433 (8.5)/	338 (8·0)/	70 (6.5)/	16 (6.6)/
D.1 (0)	2548 (11.8)***	2417 (11-2)	1657 (10-2)***	967 (8.9)	443 (8.4)
Palpitations, n (%)					
Almost never and Sometimes	327 (6.4)/	288 (5.6)/	196 (4.6)/	40 (3.7)/	1-4/
	1057 (4.9)	909 (4.2)	528 (3.2)	279 (2.6)	124 (2.4)
Often and Almost always	155 (3.0)/	145 (2.8)/	104 (2.4)/	22 (2.0)/	1-4/
	632 (2.9)	573 (2.6)	362 (2.2)	203 (1.9)	91 (1.7)
Trouble remembering or concentrating, n					
(%)					
Almost never and Sometimes	665 (13.0)/	601 (11.8)/	403 (9.5)/	81 (7.5)/	15 (6·2)/
	2594 (12.0)	2186 (10·1)	1203 (7.4)	607 (5.6)	270 (5·1)
Often and Almost always	616 (12·1)/	594 (11.6)/	457 (10.8)/	108 (10.0)/	20 (8·3)/
	2636 (12·2)	2495 (11.5)	1706 (10.5)	948 (8.7)	426 (8·1)
Cold hands/feet, n (%)					
Almost never and Sometimes	255 (5.0)/	236 (4.6)/	176 (4·1)/	39 (3.6)/	9 (3.7)/
	1022 (4.7)	875 (4.0)	494 (3.0)	279 (2.6)	129 (2.5)
Often and Almost always	406 (8.0)/	392 (7.7)/	300 (7.1)/	64 (5.9)/	11 (4.5)/
	2064 (9.5)**	1968 (9.1)	1321 (8·1)*	784 (7-2)	330 (6·3)
Cough, n (%)		, ,	\		/
Almost never and Sometimes	245 (4.8)/	209 (4.1)/	142 (3·3)/	36 (3·3)/	7 (2.9)/
	669 (3.1)	509 (2.4)	294 (1.8)	179 (1.6)	84 (1.6)
Often and Almost always	99 (1.9)/	85 (1.7)/	62 (1.5)/	13 (1.2)/	1-4/
Onen and Annost arways	245 (1.1)***	198 (0.9)	116 (0.7)***	67 (0.6)	24 (0.5)
Channel line a (0/)	243 (1.1)	170 (0.2)	110 (0.1)	07 (0.0)	24 (0.3)
Chapped lips, n (%)					

Almost never and Sometimes	229 (4.5)/	205 (4.0)/	140 (3.3)/	28 (2.6)/	1-4/
	1066 (4.9)	826 (3.8)	425 (2.6)	220 (2.0)	97 (1.8)
Often and Almost always	247 (4.8)/	234 (4.6)/	162 (3.8)/	38 (3.5)/	8 (3.3)/
·	1355 (6.3)**	1216 (5.6)	756 (4.7)*	422 (3.9)	167 (3.2)
Dizziness when standing, n (%)					
Almost never and Sometimes	528 (10·3)/	481 (9.4)/	337 (7.9)/	69 (6.4)/	17 (7.0)/
	1868 (8.6)	1604 (7.4)	931 (5.7)	501 (4.6)	232 (4.4)
Often and Almost always	348 (6.8)/	325 (6.4)/	239 (5.6)/	47 (4.3)/	5 (2.1)/
·	1387 (6.4)	1281 (5.9)	821 (5.1)	433 (4.0)	187 (3.6)
Light sensitivity, n (%)					
Almost never and Sometimes	217 (4.2)/	191 (3.7)/	130 (3.1)/	25 (2.3)/	6 (2.5)/
	854 (3.9)	718 (3.3)	414 (2.5)	232 (2·1)	114 (2.2)
Often and Almost always	175 (3.4)/	167 (3.3)/	118 (2.8)/	28 (2.6)/	5 (2·1)/
*	885 (4·1)*	830 (3.8)	578 (3.6)*	356 (3.3)	152 (2.9)
Discoloured fingers/toes, n (%)					
Almost never and Sometimes	13 (0.3)/	10 (0.2)/	8 (0.2)/	1-4/	0 (0.0)/
	146 (0.7)	122 (0.6)	71 (0.4)	45 (0.4)	18 (0.3)
Often and Almost always	16 (0.3)/	14 (0.3)/	12 (0.3)/	1-4/	1-4/
-	111 (0.5)	102 (0.5)	63 (0.4)	39 (0.4)	19 (0.4)
Extreme paleness, n (%)					
Almost never and Sometimes	45 (0.9)/	41 (0.8)/	27 (0.6)/	5 (0.5)/	1-4/
	282 (1.3)	236 (1.1)	125 (0.8)	79 (0.7)	29 (0.6)
Often and Almost always	32 (0.6)/	29 (0.6)/	21 (0.5)/	5 (0.5)/	0 (0.0)/
-	220 (1.0)*	210 (1.0)	150 (0.9)*	91 (0.8)	39 (0.7)

^a Differences between groups are tested at 2- and 6-months using Chi-Square tests.

^{* &}lt;0.05 **<0.001

^{***&}lt;0.0001

Table S2 - Psychological and social symptoms in cases and controls $% \left\{ 1,2,...,2,...\right\}$

	COVID population	Controls	p-value ^a
	(n=6630)	(n=21640)	
Being scared, n (%)			
Almost never and Sometimes	1819 (27-4)	7771 (35.9)	<0.0001
Often and Almost always	301 (4.5)	1441 (6.7)	
Feeling sad, n (%)			
Almost never and Sometimes	3607 (54·4)	13105 (60.6)	<0.0001
Often and Almost always	873 (13·2)	3748 (17-3)	
Trouble sleeping, n (%)			
Almost never and Sometimes	2852 (43.0)	10496 (48·5)	<0.0001
Often and Almost always	1199 (18·1)	4475 (20.7)	
Feeling worried about what will happen to me, n $(\%)$			
Almost never and Sometimes	1932 (29·1)	8195 (37-9)	
Often and Almost always	567 (8.6)	3321 (15·4)	<0.0001
Trouble being with peers, n (%)			
Almost never and Sometimes	1806 (27·2)	8057 (37-2)	<0.0001
Often and Almost always	256 (3.9)	1562 (7-2)	
Others don't want to be friends, n (%)			
Almost never and Sometimes	1106 (16·7)	5685 (26.3)	<0.0001
Often and Almost always	112 (1.7)	618 (2.9)	
Being teased, n (%)			
Almost never and Sometimes	489 (7.4)	2508 (11.6)	<0.0001
Often and Almost always	26 (0.4)	111 (0.5)	

^a Differences between groups were tested using Chi-Square test.

Table S3 - Sick days during the past 12 months

	COVID-population (n=6630)	Controls (n=21640)	p-value ^a	Long COVID group ^b (n=2997)
Sick days past 12 months, n (%)				
0-5 days	2684 (40.5)	13026 (60·2)	<0.0001	888 (29-6)
6-10 days	1670 (25·2)	4240 (19·6)		760 (25·4)
11-15 days	1071 (16·2)	1856 (8.6)		565 (18-9)
16 days-1 month	1078 (16·3)	1979 (9.2)		688 (23.0)
More than 1 month	127 (1.9)	539 (2.5)		96 (3·2)
Absence from school because of illness, n (%)				
0-5 days	3817 (57-6)	15732 (72.7)	<0.0001	1601 (53·4)
6-10 days	1274 (19·2)	2911 (13·5)		745 (24.7)
11-15 days	844 (12·7)	1220 (5.6)		386 (12.9)
16 days-1 month	585 (8.8)	1230 (5.7)		210 (7.0)
More than 1 month	110 (1.7)	547 (2.5)		55 (1.8)
Sick days perceived to be related to COVID-19, n (%)				
0-5 days	4222 (63·7)	NA	NA	1476 (49·3)
6-10 days	1415 (21·3)	NA		640 (21·4)
11-15 days	650 (9.8)	NA		441 (14·7)
16 days-1 month	281 (4·2)	NA		375 (12·5)
More than 1 month	62 (0.9)	NA		65 (2·2)

^a Differences between groups tested using Chi-Square test.

^b The long COVID population is defined by cases reporting at least one new-onset symptom not known before the positive SARS-CoV-2 test and present 8 weeks after the test.

 $Table \ S4-Question naire \ Long COVIDK ids DK$

Questionnaire - LongCOVIDKidsDK	
Subject	Items
General health	Weight Height Weight loss Unintended weight gain Comorbidity - Asthma - Allergy - Eczema - Tics - ADHD/ADS - Epstein Bar Virus - Arthritis - ME/CFC - Autism - Hyper-mobility - Human Papilloma Virus - OCD/anxiety/depression
SARS CoV-2 infection (cases only)	Month and year of positive SARS CoV-2 test Severity of perceived acute COVID-19 symptom burden - No symptoms - Mild symptoms - Severe symptoms
Sick days	Number of sick days the past 12 months Number of sick days the past 12 months related to SARS CoV-2 (cases only) Absence from school because of illness
CSSI	Headache Fainting or dizziness Pain in your heart or chest Feeling low in energy or slowed down Pains in your lower back Sore muscles Trouble getting your breath (when you're not exercising) Hot or cold spells (suddenly feeling hot or cold for no reason) Numbness or tingling in parts of your body Weakness (feeling weak) in parts of your body Heavy feelings in your arms or legs Nausea or upset stomach (feeling like you might throw up, or having an upset stomach) Constipation (when it's hard to have a B.M. or go poop) Loose (runny) BM's or diarrhea Pain in your stomach or abdomen (stomach aches) Your heart beating too fast (even when you're not exercising) Difficulty swallowing Losing your voice Blurred vision (when things look blurry, even with glasses on) Vomiting (or throwing up) Feeling bloated or gassy Food making you sick Pain in your knees, elbows or other joints
PedsQL	It is hard for me to walk more than one block/or one meter It is hard for me to run It is hard for me to do sports activity or exercise It is hard for me to lift something heavy It is hard for me to take a bath or shower by myself It is hard to me to do chores around the house I hurt or ache I have low energy I feel afraid or scared I feel sad or blue I feel angry I have trouble sleeping I worry about what will happen to me I have trouble getting along with other kids

	Other kids do not want to be my friend Other kids tease me I cannot do things that other kids my age can do It is hard to keep up when I play with other kids It is hard to pay attention in class I forget things I have trouble keeping up with my schoolwork I miss school because of not feeling well I miss school to go to the doctor or hospital
Long COVID symptoms	Abdominal pain Chest pain Headaches Fatigue
For how long have you been affected by the symptom?	Pain and weakness in muscle/joints Sore throat Dizziness
Are you still affected by the symptom? (cases only)	Skin rash Mood swings
Were you affected by this symptom before the	Nausea
COVID-19 infection? (cases only)	Fever
` "	Loss of appetite
Do you suspect the symptom to be related to the	Breathing problems
COVID-19 infection? (cases only)	Dark circles under the eyes
	Heart beating
	Memory or concentration difficulties
	Cold feet/hands
	Cough
	Dry and cracked lips
	Dizziness when standing Light sensitivity
	COVID fingers/toes (discoloration of fingers and toes)
	Extreme paleness

Table S5 - Covid-19 Denmark: School/national lockdown and reopening, and changes in testing policies

2020	
March 11	Lockdown: All schools, day care centers and institutions are closed.
April 15	Reopening of schools: 0-5th grade, day care centers and institutions.
May 18	Reopening of schools: 6th-10th grade and boarding schools.
May 27	Reopening of high schools and secondary educations.
December 9	Home/online schooling of pupils for 5 th to 10 th grade, high school students, and students at secondary educations.
December 21	Home/online schooling of all pupils.
2021	
February 8	Reopening of schools for 0-4 th grade
March 1	Reopening of all schools and boarding schools on the remote Danish island, Bornholm. All boarding school pupils and pupils in 9 th and 10 th grade in Northern and Western Jutland can return to school.
March 15	All boarding school pupils can return to school, pupils in 5th-8th grade attend school once a week (outdoors) and online schooling the remaining time. All 9th and 10th grade pupils, high school pupils and graduating students in secondary education, except those who live in areas around Copenhagen, can attend school part-time (50%). On small Danish islands, all pupils can return to school.
April 6	Pupils in 5 th -8 th grade return to school part-time (50%). The same applies for non-graduating students in high schools and secondary educations.
April 21	Pupils in 5 th -8 th have the opportunity for outdoor attendance during the weeks when there is no access to indoor teaching. Graduating students in primary schools and in upper secondary and adult education can attend 80% while students in secondary educations can attend fully. The reopening is geographically differentiated and covers all parts of the Denmark with exception of the metropolitan area.
May 6	Pupils in 5 th -8 th grade can return to school 100%
May 21	All students in high schools and secondary educations can return 100%
	wn and reopening
2020 March 11	Leal daying All airil converts in non-critical functions are cart home. Assembly, her for more than 100 people
Maich 11	Lockdown: All civil servants in non-critical functions are sent home. Assembly ban for more than 100 people. The government of Denmark closes its borders to all but Danish citizens, individuals with Danish work or residency permit
March 14	and those the Danish government determines have a critical purpose of travel.
March 18	Assembly for more than 10 people, closure of liberal professions, indoor sports and sports facilities, fitness centers, malls, cultural life.
April 20	Reopening of liberal professions.
May 7	Reopening of outdoor sports and sports facilities.
May 11	Reopening of all retail.
May 18	Reopening of restaurants and churches.
May 21	Reopening of museums, theaters, cinemas and zoos.
May 27	Reopening of adult education and public research activities.
June 8	Assembly ceiling is raised to 50 persons. Reopening of fitness centers and indoor swimming facilities.
July 4	Assembly ceiling is raised to 100 persons.
September 19	Assembly ceiling is lowered to 50 people, venues can still have up to 500 seated guests. Cafes, bars, and restaurants must close at 22.00. For all other workplaces - both private and public – the government encourages employees to work from home to the extent possible and appropriate.
December 17	All major shopping centers in the country close down.
December 21	All leisure activities and liberal professions, such as hairdressers, close down.
December 25	All stores except pharmacies and supermarkets close down.
2021	
2021	
March 1	Denmark partially reopens: Department stores under 5000 m2 (with the exception of department stores in malls) are reopening with stricter square meter requirements. Stores of more than 5000 m2 can be open to a very limited number of systemers by appointment. Outdoor sultural institutions reopened (requirement for pageting COVID 10 test). The assembly
	customers by appointment. Outdoor cultural institutions reopened (requirement for negative COVID-19 test). The assembly ban is raised to 25 people for outdoor sports and association activities.
March 22	The parliament has agreed to reopen society - with a few exceptions - when citizens older than 50 years have been vaccinated.
April 6	Liberal professions reopen on display of EU COVID certificate.
	Shopping malls, department stores, as well as outdoor seating and organized sports for children. Outdoor dining at
April 21	restaurants and cafes, museums, art galleries and libraries reopen on display of EU COVID certificate.
May 6	The assembly ceiling is raised to 25 indoors, venues can have up to 2,000 seated guests. Indoor dining at restaurants and cafes, conferences, venues, theatres and cinemas on presentation of EU COVID certificate. Organized indoor sports for adults over the age of 18 on presentation of EU COVID certificate.
	All remaining sports, leisure and association activities are open upon presentation of EU COVID certificate. Indoor facilities

etc.
tal,
:
onse
eive
re

 $^{* \}underline{https://askabout.io/covid-19/ask/what-is-the-government-response-timeline-for-denmark/}$